

DRAFT

Packing paths of length at least two

Mikio Kano¹ Gyula Y. Katona^{1,2} Zoltán Király^{3,*}

¹Department of Computer and Information Sciences,
Ibaraki University,
Hitachi, 316-8511 Japan

²Department of Computer and Information Sciences,
Technical University of Budapest,
H-1111 Hungary

³Department of Computer Science, Eötvös L. University, Budapest,
Rákóczi út 5. H-1088 Hungary
e-mail: kiraly@cs.elte.hu

Keywords: Path factor, graph factor, packing

Abstract

We give a simple proof for Kaneko's theorem which gives a sufficient and necessary condition for the existence of a $\{P_3, P_4, P_5\}$ -factor in a graph. Moreover we generalize this theorem and give a formula for the order of a maximum $\{P_3, P_4, P_5\}$ -packing.

1 Introduction

We consider finite graphs without multiple edges and loops. Let P_n denote the path which contains n vertices and $n - 1$ edges. For a subset X of vertices of graph G , $\langle X \rangle_G$ denotes the subgraph of G induced by X .

For a set $\{A, B, C, \dots\}$ of connected graphs, a subgraph F of a graph G is called an $\{A, B, C, \dots\}$ -packing of G if each component of F is isomorphic to one of $\{A, B, C, \dots\}$. An $\{A, B, C, \dots\}$ -packing is said to be maximum iff it covers a maximum number of vertices of G . If F is a spanning subgraph, then it is called a

*Research supported by OTKA grants OTKA T 029772 and T 030059.

perfect $\{A, B, C, \dots\}$ -packing or an $\{A, B, C, \dots\}$ -factor. Observe that a graph has a $\{P_3, P_4, P_5\}$ -factor if and only if it has a $\{P_n \mid n \geq 3\}$ -factor, which we abbreviate as $\{P_{\geq 3}\}$ -factor. We will use this evidence throughout the paper.

A graph H is said to be *factor-critical* if $H - \{v\}$ has a 1-factor for all $v \in V(H)$. For a factor-critical graph H with $V(H) = \{v_1, v_2, \dots, v_n\}$, add new vertices $\{u_1, u_2, \dots, u_n\}$ together with new edges $\{v_i u_i \mid 1 \leq i \leq n\}$ to H . Then the resulting graph is called a *sun*. Note that K_2 is a sun and by definition, we regard K_1 also as a sun (see Figure 1). We call a sun with one vertex a *small sun*, otherwise a *big sun*. We denote by $\text{Sun}(G)$ the set of sun components of G and let $\text{sun}(G) = |\text{Sun}(G)|$ the number of sun components.




Figure 1: Suns

A vertex of degree one is called a *pendant vertex*, and an edge incident with a pendant vertex is called a *pendant edge*. For a vertex v of a graph G , we denote by $\deg_G(v)$ the *degree* of v in G , and by $N_G(v)$ the *neighborhood* of v in G . For a subset $S \subseteq V(G)$, we define $N_G(S) := \cup_{x \in S} N_G(x)$.

H. Wang [10] characterized the *bipartite* graphs having a $\{P_{\geq 3}\}$ -factor. A. Kaneko recently generalized this theorem to general graphs. There are many results on component factors (for example, see [1] and [6]), but besides the well known theorem of Tutte [9] about f -factors and the more general theorem of Lovász [7] about (g, f) -factors all previous positive results (ie. that gives a good characterization) allows P_2 as a component. Hell and Kirkpatrick [4] proved that if H is a connected graph on at least 3 vertices then deciding whether a given graph G contains an $\{H\}$ -factor is *NP*-complete. Thus, for example, we do not have a good characterization of graphs having a $\{P_3\}$ -factor.

On the other hand we should mention the corresponding theorems of Hartvigsen (see [2] and [3]) about cycle-factors without short cycles.

Theorem 1 (Kaneko[5]) *A graph G has a $\{P_{\geq 3}\}$ -factor if and only if*

$$\text{sun}(G - S) \leq 2|S| \quad \text{for all } S \subset V(G). \quad (1)$$

2 A Simple Proof of Theorem 1

The following lemma is an easy consequence of Hall's theorem ([8], Theorem 1.1.3).

Lemma 2 *Let B be a bipartite graph with partite sets X and Y such that $|Y| = 2|X|$. B has a $\{P_3\}$ -factor, ie. a factor H such that $\deg_H(x) = 2$ for all $x \in X$ and $\deg_H(y) = 1$ for all $y \in Y$ if and only if*

$$|N_B(S)| \geq 2|S| \text{ for all } S \subseteq X.$$

Important properties of suns are described in the following two lemmas.

Lemma 3 *Let D be a sun which is not a K_1 , and let $u_i v_i$ be a pendant edge of D . Then $D - \{u_i, v_i\}$ has a $\{P_4\}$ -factor.*

Lemma 4 *Let D be a sun on more than two vertices, and v' is a pendant vertex of D . $D - v'$ has a $\{P_4, P_5\}$ -factor.*

PROOF: Let $V(D) = \{v, x, y, z, \dots\} \cup \{v', x', y', z', \dots\}$, where y' is the pendant vertex connected to y etc. Let F be the factor-critical graph $\langle \{v, x, y, z, \dots\} \rangle_D$. Choose a neighbor x of v in F . Now $F - x$ has a perfect matching M , with some $vy \in M$. Take the path $\{y', y, v, x, x'\}$ and the $\{P_4\}$ -s extending the other edges of M by pendant ones. \square

PROOF OF THEOREM 1: Since no sun component can have a $\{P_{\geq 3}\}$ -factor, it is easy to show that if G has a $\{P_{\geq 3}\}$ -factor, then (1) holds.

We now prove the sufficiency by induction on $\|G\| = |E(G)|$. Suppose that G satisfies (1). By setting $S = \emptyset$ condition (1) implies that no component of G is a sun. We may assume that G is connected and $|G| \geq 3$. We consider some cases.

Case 1 There exists $\emptyset \neq S \subset V(G)$ such that $\text{sun}(G - S) = 2|S|$.

Choose a subset S of $V(G)$ so that S is maximal among all the subsets S' satisfying $\text{sun}(G - S') = 2|S'|$.

Let C be any non-sun component of $G - S$. Then for a subset $X \subset V(C)$, we have

$$2|S \cup X| \geq \text{sun}(G - (S \cup X)) = \text{sun}(G - S) + \text{sun}(C - X) = 2|S| + \text{sun}(C - X).$$

Thus $\text{sun}(C - X) \leq 2|X|$. Hence C satisfies (1), and so C has a $\{P_{\geq 3}\}$ -factor by induction.

We define the bipartite graph B with vertex set $S \cup \text{Sun}(G - S)$ by contracting every sun-component into a single vertex and removing multiple edges and edges inside S . Now $|\text{Sun}(G - S)| = 2|S|$, and we show that

$$|N_B(X)| \geq 2|X| \text{ for all } X \subseteq S. \tag{2}$$

Suppose that $|N_B(Y)| < 2|Y|$ holds for some $Y \subseteq S$. Then $\text{Sun}(G - (S \setminus Y)) \supseteq \text{Sun}(G - S) \setminus N_B(Y)$ holds, and thus

$$\text{sun}(G - (S \setminus Y)) \geq \text{sun}(G - S) - |N_B(Y)| > 2|S| - 2|Y| = 2|S \setminus Y|,$$

is implied, which contradicts the assumption (1). Thus (2) holds.

Therefore, by Lemma 2 graph B has a factor H such that $\deg_H(x) = 2$ for all $x \in S$ and $\deg_H(C) = 1$ for all $C \in \text{Sun}(G - S)$. By making use of this factor, by Lemma 3 we can obtain a $\{P_{\geq 3}\}$ -factor of G (see Figure 2).




Figure 2: Extension of the $\{P_3\}$ -factor of B to a $\{P_{\geq 3}\}$ -factor in G

Case 2 $\text{sun}(G - S) < 2|S|$ for all $\emptyset \neq S \subset V(G)$ and there exists $\emptyset \neq S' \subset V(G)$ for which $\text{sun}(G - S') = 2|S'| - 1$.

Choose a subset S so that S is maximal among all subsets S' satisfying $\text{sun}(G - S') = 2|S'| - 1$.

Let C be any non-sun component of $G - S$ and let $\emptyset \neq X \subset V(C)$. Using the maximality of S we obtain

$$2|S \cup X| - 2 \geq \text{sun}(G - (S \cup X)) = \text{sun}(G - S) + \text{sun}(C - X) = 2|S| - 1 + \text{sun}(C - X).$$

$$\text{Thus } \text{sun}(C - X) \leq 2|X| - 1. \quad (3)$$

Hence C has a $\{P_{\geq 3}\}$ -factor by induction.

Claim 1 *If $G - S$ has a non-sun component then the desired $\{P_{\geq 3}\}$ -factor exists.*

PROOF: Let C be such a component, $v \in S$ and $w \in C$ such that vw is an edge. Let w^* be a new vertex and consider the graph $H := \langle C \rangle_G + ww^*$. Using (3) it is easy to see that H satisfies (1) for nonempty sets: $\text{sun}(H - X) \leq \text{sun}(C - X) + 1 \leq 2|X| - 1 + 1$. Clearly $\|H\| < \|G\|$, so by induction H has a $\{P_{\geq 3}\}$ -factor containing a path P ending with $\{w, w^*\}$ or H itself is a sun. In the latter case, by Lemma 3, H has a $\{P_2, P_4\}$ -factor so that the only P_2 is $P = \{w, w^*\}$.

Construct bigraph B as above, but add a new pendant edge vw' to it. Now B satisfies (2) and we obtain a $\{P_3\}$ -factor containing a path Q ending with $\{v, w'\}$. Now take $P - w^*$ and $Q - w'$ and join them by the edge vw . Using this factor we can obtain a $\{P_{\geq 3}\}$ -factor in the same way as in the previous case. \square

Claim 2 *If there exists $v \in S$ connected to zero, or more than one small suns in $\text{Sun}(G - S)$ then the desired $\{P_{\geq 3}\}$ -factor exists.*

PROOF: Construct B as before with the additional pendant edge vw' . Extend the $\{P_3\}$ -factor of B as before to obtain a $\{P_{\geq 3}\}$ -factor of $G + w'$ and delete now w' . The path containing v becomes a path of length one $\{v, w\}$ if $\{w\}$ is a small sun connected to v . By our assumptions another small sun w^* is also connected to v , and w^* is an endvertex of another path that can be extended by adding edge w^*v . \square

Claim 3 *If $\{w\} \in \text{Sun}(G - S)$ a small sun and w is not pendant in G then the desired $\{P_{\geq 3}\}$ -factor exists.*

PROOF: As w is not pendant, it is connected to some $v \in S$ and $v' \in S$, $v' \neq v$. Construct B as before with additional pendant edge vw' . We claim that $B' = B - vw$ satisfies (2). If $X \subset S$ then either $v \notin X$ and $N_{B'}(X) = N_B(X)$ or $v \in X$ and $|N_{B'}(X)| \geq 2|X| + 1$ otherwise $S \setminus X$ would be a set with $\text{sun}(G - (S \setminus X)) \geq 2|S \setminus X|$. For $X = S$ we need to prove $w \in N_{B'}(S)$ which is true because wv' is an edge. Now take the path ending in w' in the $\{P_{\geq 3}\}$ -factor obtained using the $\{P_3\}$ -factor of B' , delete w' and connect the remains of this path to the path ending in w by edge vw . \square

So we may assume from now on that there are $|S|$ small suns in $\text{Sun}(G - S)$ and $S - 1$ big suns. Each small sun is a pendant vertex of G and they are connected to different vertices in S .

Claim 4 *If every vertex of G with degree ≥ 2 has a pendant neighbor then the desired $\{P_{\geq 3}\}$ -factor exists.*

PROOF: Let U be the set of vertices with degree ≥ 2 . If $\langle U \rangle_G$ has a perfect matching, we are done. Otherwise there exists $X \subset U$ such that there are more than $|X|$ factor-critical components in $\langle U \setminus X \rangle_G$, consequently $\text{sun}(G - X) > 2|X|$, which is a contradiction. \square

So we may assume there is a vertex with degree ≥ 2 which has no pendant neighbor in G . Clearly it is in a big sun $D \in \text{Sun}(G - S)$. This means that $\langle D \rangle_G$

has a pendant vertex v' with neighbor $v \in D$ so that v' is connected (in G) to some $u \in S$ and if D is a K_2 -component then v is also connected to S .

Subcase 2.1 $|D| = 2$

Let G' be the graph obtained from G by deleting the edge vv' . Now $\text{sun}(G' - S) = 2|S|$. Construct B as in Case 1. It is easy to see that (2) is satisfied, so we obtain a $\{P_{\geq 3}\}$ -factor as in Case 1.

Subcase 2.2 $|D| > 2$

Let the small sun neighbor of u be $D = \{w\}$. Construct B as before by adding pendant edge uw' . Take the $\{P_3\}$ -factor of B , construct a $\{P_{\geq 3}\}$ -factor of $G + w'$ and delete w' . Now we have a $\{P_{\geq 3}\}$ -factor of G unless $\{w, u\}$ is a path of length one in this factor. If v' is an endvertex of another path of this factor then we can extend $\{w, u\}$ by adding edge uv' . Otherwise the path containing v' ends with $\{s, v', v\}$ (by our construction this is the only possibility) where $s \in S, s \neq u$. Now delete edge $v'v$ from the factor as well as the edges inside D , add edge uv' , and use Lemma 4 to obtain a $\{P_{\geq 3}\}$ -factor of $D - v'$.

Case 3 $\text{sun}(G - S) \leq 2|S| - 2$ for all $\emptyset \neq S \subset V(G)$.

If G has a pendant vertex u connected to v , then $\text{sun}(G - \{v\}) \geq 1$, which contradicts the assumption of this case. Thus G is not a tree, and so we can find an edge e for which $G - e$ is connected. For every subset $\emptyset \neq S \subset V(G - e)$, we have

$$\text{sun}((G - e) - S) \leq \text{sun}(G - S) + 2 \leq 2|S| - 2 + 2 = 2|S|.$$

Moreover, $G - e$ is not a sun because having at least three vertices it would be a sun with at least three pendant vertices, but in this case G would have at least one pendant vertex as well. Therefore, $G - e$ has a $\{P_{\geq 3}\}$ -factor by the inductive hypothesis, and so does G .

Consequently, the proof is complete. \square

3 The Order of a Maximum $\{P_{\geq 3}\}$ -packing

Lemma 5 *Let B be a bipartite graph with bipartition $X \cup Y$ and $Y^* \subseteq Y$. Define*

$$\text{def}(B) := \max_{Y' \subseteq Y} (|Y'| - 2|N_B(Y')|)$$

and

$$\text{def}^*(B) := \max_{Y' \subseteq Y^*} (|Y'| - 2|N_B(Y')|).$$

If $\text{def}(B) = |Y| - 2|X|$ then B has a $\{P_3\}$ -packing which covers $|Y| - \text{def}(B) = 2|X|$ vertices of Y including $|Y^| - \text{def}^*(B)$ vertices of Y^* , and covers all vertices in X .*

PROOF: Define bipartite graph $B' = ((X \cup X') \cup Y, E')$ by adding for every vertex $x \in X$ a new vertex $x' \in X'$ and connecting x' to all neighbors of x . By Ore's theorem ([8], Theorem 1.3.1) there exist a matching M that covers $|Y| - \text{def}(B') = |X \cup X'|$ vertices of Y and (consequently) all vertices of $X \cup X'$. Moreover, there exists another matching M^* that covers $|Y^*| - \text{def}^*(B')$ vertices of Y^* . It is well known that this implies the existence of a matching which covers $X \cup X'$ and $|Y^*| - \text{def}^*(B')$ vertices of Y^* (see [8], Exercise 1.4.3). This gives the desired $\{P_3\}$ -packing in B if we contract all pairs x, x' . \square

Let $k_2(H)$ denote the number of components of H which consist of an edge, in other terms the number of sun components isomorphic to a K_2 .

Theorem 6 *The order of a maximum $\{P_{\geq 3}\}$ -packing in a graph G is*

$$\text{pp}(G) := |V(G)| - \max_{T \subseteq S \subseteq V(G)} (\text{sun}(G - S) - 2|S| + k_2(G - T) - 2|T|).$$

PROOF: It is proved first that the above expression is an upper bound on the order of a maximum $\{P_{\geq 3}\}$ -packing. Let $T \subseteq S \subseteq V(G)$ such that $|V(G)| - \text{pp}(G) = \text{sun}(G - S) - 2|S| + k_2(G - T) - 2|T|$. Clearly if F is a $\{P_{\geq 3}\}$ -packing of G then there is a vertex in at least $\text{sun}(G - S) - 2|S|$ components in $\text{Sun}(G - S)$ which cannot be covered by F . Moreover, there are at least $k_2(G - T) - 2|T|$ K_2 -components of $G - T$ where none of the two vertices can be covered.

To prove the other direction choose $T \subseteq S \subseteq V(G)$ such that $\text{sun}(G - S) - 2|S| + k_2(G - T) - 2|T|$ is maximum.

Let C be a non-sun component of $G - S$. Then for a subset $X \subseteq V(C)$ we have $\text{sun}(G - (S \cup X)) - 2|S \cup X| + k_2(G - T) - 2|T| \leq \text{sun}(G - S) - 2|S| + k_2(G - T) - 2|T|$

by the choice of S and T . Since $\text{sun}(G - (S \cup X)) = \text{sun}(G - S) + \text{sun}(C - X)$ and $X \cap S = \emptyset$, it follows that $\text{sun}(C - X) \leq 2|X|$ which implies that C has a $\{P_{\geq 3}\}$ -factor. Hence, we may assume from now on that $G - S$ has only sun components.

Construct a bipartite graph B from G by contracting each sun component into a single vertex and removing multiple edges and edges inside S . The set of vertices which arose from the sun components in $G - S$ is denoted by Y , the set of vertices that arose from the contraction of K_2 components of $G - T$ is denoted by Q and the set of vertices which arose from the contraction of K_2 components of $G - S$ is denoted by Y^* . The K_2 components of $G - T$ are elements of $\text{Sun}(G - S)$, because if there exists a K_2 component of $G - T$ such that $V(D) \cap (S - T) \neq \emptyset$ then we get a contradiction by considering $S \setminus V(D)$ and T . Thus $Q \subseteq Y^* \subseteq Y$ holds.

First we show that $\text{def}^*(B) = k_2(G - T) - 2|T|$. Suppose that

$$R - 2|N_B(R)| > k_2(G - T) - 2|T|$$

holds for some $R \subseteq Y^*$. Then choosing $N_B(R)$ instead of T (and keeping S), obviously violates the choice of T and S .

Next we prove that $\text{def}(B) = \text{sun}(G - S) - 2|S|$. Suppose that

$$R - 2|N_B(R)| > \text{sun}(G - S) - 2|S|$$

holds for some $R \subseteq Y$. Let $S' := N_B(R \cup Q)$ and $T' := N_B(R \cap Q)$. Since $N_B(Q) = T$, it is obvious that $S' = N_B(R) \cup T$ and $T' \subseteq N_B(R) \cap T$. In this way

$$\begin{aligned} \text{sun}(G - S') - 2|S'| + k_2(G - T') - 2|T'| &\geq |R \cup Q| - 2|S'| + |R \cap Q| - 2|T'| = \\ &|R| + |Q| - 2|S'| - 2|T'| \geq |R| + |Q| - 2|N_B(R) \cup T| - 2|N_B(R) \cap T| = \\ &|R| + |Q| - 2|N_B(R)| - 2|T| > \text{sun}(G - S) - 2|S| + k_2(G - T) - 2|T| \end{aligned}$$

holds, which contradicts the choice of S and T .

By applying Lemma 5 a $\{P_3\}$ -packing is obtained in B which covers all vertices in S , $|Y| - \text{sun}(G - S) + 2|S|$ vertices of Y including $|Y^*| - k_2(G - T) + 2|T|$ vertices of Y^* . (See Figure 3.)




Figure 3: B with the desired factor.

In the original graph G we can extend this packing in the usual way by Lemma 3 (see Figure 2). On the other hand, the sun-components with more than two vertices which is not covered by the P_3 -s in B can be almost covered by a $\{P_{\geq 3}\}$ -packing by Lemma 4. This gives a $\{P_{\geq 3}\}$ -packing of size $\text{pp}(G)$. \square

Note, that

$$\max_{S \subseteq V(G)} (\text{sun}(G - S) - 2|S|) = 0 \quad (4)$$

implies that $k_2(G - T) - 2|T| \leq 0$ holds for any $T \subseteq V(G)$, because K_2 is a sun. This shows that

$$\max_{T \subseteq S \subseteq V(G)} (\text{sun}(G - S) - 2|S| + k_2(G - T) - 2|T|) = 0$$

holds if and only if (4) holds, hence Theorem 6 implies Theorem 1.

4 Acknowledgement

The authors are indebted to the organizers and sponsors of the 1st Japanese-Hungarian Symposium on Discrete Mathematics and Its Applications (1999, Kyoto, Japan), where they learned the result of Kaneko and started to work on this paper.

References

- [1] J. Akiyama and M. Kano, Factors and factorizations of graphs – A survey, *J. Graph Theory*, **9** (1985) 1-42.
- [2] D. Hartvigsen, *Extensions of Matching Theory*, PhD. Thesis, Carnegie-Mellon University (1984).
- [3] D. Hartvigsen, *The Square-Free 2-Factor Problem in Bipartite Graphs*, Integer Programming and Combinatorial Optimization, 7th IPCO Conference (Graz, 1999) LNCS 1610, 234-241.
- [4] P. Hell and D. G. Kirkpatrick, *On the completeness of a generalized matching problem*, Proceedings of the Tenth Annual ACM Symposium on Theory of Computing (San Diego, 1978), 240-245.
- [5] A. Kaneko, A necessary and sufficient condition for the existence of a path factor every component of which is a path of length at least two, submitted.
- [6] M. Loeb1 and S. Poljak, Subgraph packing – A survey, *Topics in Combinatorics and Graph Theory*, ed by R. Bodendiek and R. Henn, Physical-Verlag Heidelberg (1990).
- [7] L. Lovász, *Subgraphs with prescribed valencies*, J. Combin. Theory **8** (1970), 391-416.
- [8] L. Lovász and M. D. Plummer, *Matching Theory* (Annals of Discrete Math. 29), North-Holland (1986), Amsterdam.
- [9] W. T. Tutte, *The factors of graphs*, Canad. J. Math. **4** (1952), 314-328.
- [10] H. Wang, *Path factors of bipartite graphs*, J. Graph Theory **18** (1994), no. 2, 161-167.