

# Balancing colored points on a line by exchanging intervals

Atsushi Kaneko<sup>1</sup>, Mikio Kano<sup>2</sup> and Mamoru Watanabe<sup>3</sup>

<sup>1</sup> Koshigaya, Saitama, Japan

<sup>2</sup> Ibaraki University, Hitachi, Ibaraki, Japan

mikio.kano.math@vc.ibaraki.ac.jp

<sup>3</sup> Kurashiki, Okayama, Japan,

## Abstract

Assume that  $2a$  red points,  $2b$  blue points and  $2c$  green points lie on a line, and they are bisected into the left part  $I$  and the right part  $J$  by a point so that each of them contains  $a + b + c$  points. Then we show that there exist a point set  $X \subset I$  and a point set  $Y \subset J$  such that both  $X$  and  $Y$  consist of consecutive points,  $|X| = |Y|$ , and each of  $I - X + Y$  and  $J - Y + X$  contains exactly  $a$  red points,  $b$  blue points and  $c$  green points. Moreover we extend this result to multi-colored point sets.

## 1 Introduction

Various topics on red and blue points in the plane have been studied [1], and in the proofs of some theorems, results on colored points on a line play important role [1], [2]. In this paper, we consider some problems of 3-colored points and multi-colored points on a line.

Assume that colored  $2n$  points lie on a line, and they are bisected into the left part  $I$  and the right part  $J$  by a point so that both  $I$  and  $J$  contain precisely  $n$  points each. If the number of points of each color is even and both  $I$  and  $J$  contain the same number of points of each color, then we say that  $I$  and  $J$  are *balanced*.

In this paper, we shall prove the following three theorems, which say that the above  $I$  and  $J$  can be balanced by exchanging two subsets  $X \subset I$  and  $Y \subset J$  consisting of small number of intervals of  $I \cup J$ . Moreover, their proofs give polynomial-time algorithms for finding such subsets  $X$  and  $Y$ .

Note that if  $X$  and  $Y$  are disjoint sets, we often write  $X + Y$  for  $X \cup Y$ . Moreover, if  $Z$  is a subset of  $X$ , we often write  $X - Z$  for  $X \setminus Z$ .

**Theorem 1.** *Assume that  $2a$  red points,  $2b$  blue points and  $2c$  green points lie on a line, where  $a, b, c$  are positive integers, and they are bisected into the left part  $I$  and the right part  $J$  by a point so that each of them contains precisely  $a + b + c$  points. Then there exist a point set  $X \subset I$  and a point set  $Y \subset J$  such that both  $X$  and  $Y$  consist of consecutive points,  $|X| = |Y|$ , and both  $I - X + Y$  and  $J - Y + X$  are balanced (see Fig. 1).*




Figure 1: Red, blue and green points lying on a line, two point sets  $X \subset I$  and  $Y \subset J$ , and two balanced sets  $I - X + Y$  and  $J - Y + X$ .

If two colored points lie on a line, we can obtain a slightly stronger result as follows:




Figure 2: Red and blue points lying on a line, two point sets  $X \subset I$  and  $Y \subset J$ , and two balanced sets  $I - X + Y$  and  $J - Y + X$ .

**Theorem 2.** *Assume that  $2a$  red points and  $2b$  blue points lie on a line, where  $a$  and  $b$  are positive integers, and they are bisected into the left part  $I$*

and the right part  $J$  by a point so that each of them contains precisely  $a + b$  points. Then there exist a point set  $X \subset I$  and a point set  $Y \subset J$  such that both  $X$  and  $Y$  consist of consecutive points starting at the right end-point of  $I$  and  $J$  respectively,  $|X| = |Y|$ , and both  $I - X + Y$  and  $J - Y + X$  are balanced (see Fig. 2).

If the number of colors is more than three, we can obtain balanced sets by exchanging two or more intervals of  $I \cup J$ .




Figure 3: Four colored points lying on a line, two point sets  $X \subset I$  and  $Y \subset J$ , and two balanced sets  $I - X + Y$  and  $J - Y + X$ .

**Theorem 3.** Let  $r \geq 2$  be an integer, and let  $c_1, c_2, \dots, c_r$  be  $r$  colors. Assume that  $2n_i$  points of color  $c_i$  lie on a line for every  $1 \leq i \leq r$ . Furthermore they are bisected into the left part  $I$  and the right part  $J$  by a point so that each of them contains precisely  $n_1 + n_2 + \dots + n_r$  points. Then there exist point sets  $X \subset I$  and  $Y \subset J$  such that  $X \cup Y$  consist of at most  $\lfloor (r + 2)/2 \rfloor$  intervals of  $I \cup J$ ,  $|X| = |Y|$ , and both  $I - X + Y$  and  $J - Y + X$  are balanced (see Fig. 3). Moreover, the bound  $\lfloor (r + 2)/2 \rfloor$  is sharp.

## 2 Proofs of Theorems

For a positive integer  $d$ , we denote by  $\mathbf{R}^d$  the  $d$ -dimensional Euclidean space. Note that  $\mathbf{R}^1$  is often written  $\mathbf{R}$ . For a positive number  $\alpha$ , the curve  $\{(t, t^2, \dots, t^d) : 0 \leq t \leq \alpha\}$  in  $\mathbf{R}^d$  is called *the moment curve*. The moment curve has the following property.

**Lemma 4** (Lemma 1.6.4 of [3]). *Every hyperplane of  $\mathbf{R}^d$  intersects the moment curve in  $\mathbf{R}^d$  at most  $d$  points.*

The next theorem is well-known.

**Theorem 5** (Ham-sandwich theorem, [3], [4]). *Every  $d$  point sets in  $\mathbf{R}^d$  each of which contains even number of points can be simultaneously bisected by a hyperplane. Moreover, there is a polynomial-time algorithm for finding such a hyperplane.*

We are ready to prove theorems. We first prove Theorem 1.

*Proof of Theorem 1.* We may assume that the points of  $I \cup J$  are contained in the interval  $[0, 1]$  of  $\mathbf{R}$ . We replace the consecutive points of  $I \cup J$  along the moment curve  $\gamma = \{(t, t^2, t^3) : 0 \leq t \leq 1\}$  in the space  $\mathbf{R}^3$ . By Ham-sandwich theorem, there exists a plane  $h$  in the space that simultaneously bisects these three colored points and intersects the moment curve  $\gamma$  at most three points.

First assume that the hyperplane  $h$  intersects  $\gamma$  at three points. Let  $P_1, P_2, P_3, P_4$  denote the four sets of colored points on  $\gamma$  divided by  $h$ . By symmetry of  $I$  and  $J$ , we may assume that  $P_1 \cup P_2 \cup Q = I$  and  $R \cup P_4 = J$ , where  $Q = P_3 \cap I$  and  $R = P_3 \cap J$  and it may occur that  $Q$  or  $R$  is an empty set. Moreover, each of  $P_1 \cup P_3$  and  $P_2 \cup P_4$  contains exactly  $a$  red points,  $b$  blue points and  $c$  green points since  $h$  is a bisector.

Let  $X = P_2$  and  $Y = R$ . Then  $I - X + Y = P_1 \cup P_3$  and  $J - Y + X = P_2 \cup P_4$  are balanced. This implies  $|I| - |X| + |Y| = |J| - |Y| + |X|$ , and thus  $|X| = |Y|$ .

We next consider the case where the plane  $h$  intersects  $\gamma$  at two points. Let  $P_1, P_2, P_3$  denote the three sets of points on  $\gamma$  divided by  $h$ . By symmetry, we may assume that  $P_1 \cup Q = I$  and  $R \cup P_3 = J$ , where  $Q = P_2 \cap I$  and  $R = P_2 \cap J$ . Moreover, each of  $P_1 \cup P_3$  and  $P_2$  contains exactly  $a$  red points,  $b$  blue points and  $c$  green points since  $h$  is a bisector. Then  $I - Q + P_3 = P_1 \cup P_3$  and  $J - P_3 + Q = P_2$  are balanced, and this also implies  $|Q| = |P_3|$ . If the plane  $h$  intersects  $\gamma$  at one point, then  $I$  and  $J$  are balanced, and so the theorem holds for  $X = Y = \emptyset$ . Consequently Theorem 1 is proved.  $\square$

*Proof of Theorem 2.* We may assume that the points of  $I \cup J$  are contained in the interval  $[0, 1]$  of  $\mathbf{R}$ . We place the consecutive points of  $I \cup J$  along the the moment curve  $\gamma = \{(t, t^2) : 0 \leq t \leq 1\}$  in the plane  $\mathbf{R}^2$ . By Ham-sandwich theorem, there exists a line  $\ell$  that simultaneously bisects these red and blue points and intersects  $\gamma$  at most two points. If  $\ell$  intersects  $\gamma$  at one point, then  $I$  and  $J$  are balanced. Thus we may assume that  $\ell$  and  $\gamma$  intersect at two points.

Let  $P_1, P_2, P_3$  denote the three sets of colored points on  $\gamma$  divided by  $\ell$ . By symmetry, we may assume that  $P_1 \cup Q = I$  and  $R \cup P_3 = J$ , where  $Q = P_2 \cap I \neq \emptyset$  and  $R = P_2 \cap J \neq \emptyset$ . Moreover, each of  $P_1 \cup P_3$  and  $P_2$

contains exactly  $a$  red points and  $b$  blue points since  $\ell$  is a bisector. Hence it follows that  $I - Q + P_3 = P_1 \cup P_3$  and  $J - P_3 + Q = P_2$  are balanced, and  $|Q| = |P_3|$ . Furthermore,  $Q$  and  $P_3$  contain the right end-point of  $I$  and  $J$ , respectively. Consequently Theorem 2 is proved.  $\square$

*Proof of Theorem 3.* We may assume that the points of  $I \cup J$  are contained in the interval  $[0, 1]$  of  $\mathbf{R}$ . We place the consecutive points of  $I \cup J$  along the moment curve  $\gamma = \{(t, t^2, \dots, t^r) : 0 \leq t \leq 1\}$  in  $\mathbf{R}^r$ . By Ham-sandwich theorem, there exists a hyperplane  $h$  that simultaneously bisects these  $r$  colored points and intersects  $\gamma$  at most  $r$  points.

Let  $P_1, P_2, \dots, P_s$  denote  $s$  sets of colored points on  $\gamma$  divided by  $h$ , where  $h$  intersects  $\gamma$  at  $s-1$  points and  $2 \leq s \leq r+1$ . If  $s = 2$ , then  $P_1 = I$  and  $P_2 = J$  are balanced, and so we may assume  $3 \leq s \leq r+1$ . Since  $h$  is a bisector,  $P_1 \cup P_3 \cup \dots \cup P_s$  (or  $P_{s-1}$ ) and  $P_2 \cup P_4 \cup \dots \cup P_{s-1}$  (or  $P_s$ ) are balanced. In particular, they contain the same number of points of every color. Moreover, we can write  $I = P_1 \cup P_2 \cup \dots \cup P_{t-1} \cup Q$  and  $J = R \cup P_{t+1} \cup \dots \cup P_s$ , where  $Q = I \cap P_t$ ,  $R = J \cap P_t$ ,  $2 \leq t < s$  and it may occur that  $Q$  or  $R$  is an empty set.

We first assume that  $t$  is even. Let  $X = P_2 \cup P_4 \cup \dots \cup P_{t-2} \cup Q$  and  $Y = P_{t+1} \cup P_{t+3} \cup \dots \cup P_s$  (or  $P_{s-1}$ ). Then  $I - X + Y = P_1 \cup P_3 \cup \dots \cup P_s$  (or  $P_{s-1}$ ) and  $J - Y + X = P_2 \cup P_4 \cup \dots \cup P_{s-1}$  (or  $P_s$ ) are balanced. This implies  $|I| - |X| + |Y| = |J| - |Y| + |X|$  and thus  $|X| = |Y|$ . Moreover,  $X \cup Y$  consists of

$$\begin{aligned} \frac{t}{2} + \frac{s - (t+1)}{2} + 1 &= \frac{s+1}{2} \quad \text{or} \\ \frac{t}{2} + \frac{s-1-(t+1)}{2} + 1 &= \frac{s}{2} \end{aligned}$$

intervals. Since  $s \leq r+1$ ,  $X \cup Y$  consists of at most  $\lfloor (r+2)/2 \rfloor$  intervals.

Next assume that  $t$  is odd. Let  $X = P_2 \cup P_4 \cup \dots \cup P_{t-1}$  and  $Y = R \cup P_{t+2} \cup P_{t+4} \cup \dots \cup P_s$  (or  $P_{s-1}$ ). Then  $I - X + Y = P_1 \cup P_3 \cup \dots \cup P_s$  (or  $P_{s-1}$ ), and  $J - Y + X = P_2 \cup P_4 \cup \dots \cup P_{s-1}$  (or  $P_s$ ) are balanced. This implies  $|I| - |X| + |Y| = |J| - |Y| + |X|$  and thus  $|X| = |Y|$ . Moreover,  $X \cup Y$  consists of

$$\begin{aligned} \frac{t-1}{2} + \frac{s-t}{2} + 1 &= \frac{s+1}{2} \quad \text{or} \\ \frac{t-1}{2} + \frac{s-1-t}{2} + 1 &= \frac{s}{2} \end{aligned}$$

intervals. Since  $s \leq r+1$ ,  $X \cup Y$  consists of at most  $\lfloor (r+2)/2 \rfloor$  intervals. Consequently the existence of  $X$  and  $Y$  in Theorem 3 is proved.

We finally show that the bound  $\lfloor (r+2)/2 \rfloor$  is sharp. Consider the following colored point configuration on a line, in which the number of points of each color  $c_i, 1 \leq i \leq r-1$ , is two and the number of points of color  $c_r$  is  $2(r-1)$ . The left part  $I$  contains all the points of color  $c_i, 1 \leq i \leq r-1$ , and two points of the same color appear consecutively, and  $2(r-1)$  points of color  $c_r$  lie on the right part  $J$ . Then in order to obtain balanced sets, we have to exchange one point of each color  $c_i, 1 \leq i \leq r-1$ , of  $I$  and  $r-1$  points of color  $c_r$  of  $J$ . Therefore, if  $r$  is odd, then  $X \cup Y$  consists of at least  $(r-1)/2 + 1 = (r+1)/2$  intervals of  $I \cup J$  (see the first example of Fig. 4). If  $r$  is even, then  $X \cup Y$  consists of at least  $((r-2)/2 + 1) + 1 = (r+2)/2$  intervals of  $I \cup J$  (see the second example of Fig. 4). Hence the bound  $\lfloor (r+2)/2 \rfloor$  is sharp.  $\square$




Figure 4: Five colored points lie on a line, and  $X \cup Y$  consists of 3 intervals. Six colored points lie on a line, and  $X \cup Y$  consists of 4 intervals.

**Acknowledgement** Mikio Kano is supported by JSPS KAKENHI Grant Number 16K05248.

## References

- [1] Kaneko, A. and Kano, M.: Discrete geometry on red and blue points in the plane — A survey, *Discrete and Computational Geometry, Algorithms and Combinatorics*, Springer, Vol. 25, pp. 551-570 (2003).
- [2] Kano, M, Suzuki, K and Uno, M.: Properly colored geometric matchings and 3-trees without crossings on multicolored points in the plane, *Discrete and Computational Geometry and Graphs*, Springer, Vol. LNCS 8845, pp. 96–111 (2014).
- [3] Matoušek, J.: *Using the Borsuk-Ulam Theorem*, Springer (2003).

- [4] Lo, C-Y., Matoušek, J. and Steiger, W. L.: Algorithms for Ham-Sandwich Cuts, *Discrete and Computational Geometry*, Vol. 11, pp.433–452 (1994).